

Pattern Count on Multiply Restricted Permutations

Alina F. Y. Zhao

School of Mathematical Sciences and Institute of Mathematics

Nanjing Normal University, Nanjing 210023, PR China

alinazhao@njnu.edu.cn

Abstract

Previous work has studied the pattern count on singly restricted permutations. In this work, we focus on patterns of length 3 in multiply restricted permutations, especially for double and triple pattern-avoiding permutations. We derive explicit formulae or generating functions for various occurrences of length 3 patterns on multiply restricted permutations, as well as some combinatorial interpretations for non-trivial pattern relationships.

Mathematics Subject Classification: 05A05, 05A15, 05A19

1 Introduction

Let $\sigma = \sigma_1\sigma_2\cdots\sigma_n$ be a permutation in the symmetric group S_n written in one-line notation, and σ_i is said to be a left to right maximum (resp. right to left maximum) if $\sigma_i > \sigma_j$ for all $j < i$ (resp. $j > i$). For a permutation $q \in S_k$, we say that σ contains q as a pattern if there exist $1 \leq i_1 \leq i_2 \leq \cdots \leq i_k \leq n$ such that the entries $\sigma_{i_1}\sigma_{i_2}\cdots\sigma_{i_k}$ have the same relative order as the entries of q , i.e., $q_j < q_l$ if and only if $\sigma_{i_j} < \sigma_{i_l}$ whenever $1 \leq j, l \leq k$. We say that σ avoids q if σ does not contain q as a pattern.

For a pattern q , denote by $S_n(q)$ the set of all permutations in S_n that avoiding the pattern q , and for $R \subseteq S_k$, we denote by $S_n(R) = \bigcap_{q \in R} S_n(q)$, i.e., the set of permutations in S_n which avoid every pattern contained in R . For two permutations σ and q , we set $f_q(\sigma)$ as the number of occurrences of q in σ as a pattern, and we further denote the number of occurrences of q in a permutation set Ω by $f_q(\Omega) = \sum_{\sigma \in \Omega} f_q(\sigma)$.

Recently Cooper [5] raised the problem of determining the total number $f_q(S_n(r))$ of all q -patterns in the r -avoiding permutations of length n . Bóna [3] discovered the generating functions of the sequence $f_q(S_n(132))$ for the monotone q , and Bóna [2] further studied the generating functions for other length 3 patterns in the set $S_n(132)$,

$S_n(123, 132)$	$f_{213}(n) = (n-3)2^{n-2} + 1$	$S_n(123, 132, 213)$	$\sum f_{231}(n)x^n = \sum f_{312}(n)x^n = \frac{x^3(1+2x)}{(1-x-x^2)^3}$
	$f_{231}(n) = f_{312}(n) = (n^2 - 5n + 8)2^{n-3} - 1$		$\sum f_{321}(n)x^n = \frac{x^3(1+6x+12x^2+8x^3)}{(1-x-x^2)^4}$
	$f_{321}(n) = (n^3/3 - 2n^2 + 14n/3 - 5)2^{n-2} + 1$		$f_{213}(n) = f_{312}(n) = \binom{n}{3}$
$S_n(132, 213)$	$f_{123}(n) = (n-4)2^{n-1} + n + 2$	$S_n(123, 132, 231)$	$f_{321}(n) = (n-2)\binom{n}{3}$
	$f_{231}(n) = f_{312}(n) = (\frac{n^2}{4} - \frac{7n}{4} + 4)2^n - n - 4$		$f_{123}(n) = f_{312}(n) = \binom{n+1}{4}$
	$f_{321}(n) = (\frac{1}{12}n^3 - \frac{3}{4}n^2 + \frac{38}{12}n - 6)2^n + n + 6$		$f_{321}(n) = n(n-2)(n-1)^2/12$
$S_n(132, 231)$	$f_{123}(n) = f_{213}(n) = f_{312}(n) = f_{321}(n) = \frac{2^n}{8}\binom{n}{3}$	$S_n(123, 132, 312)$	$f_{213}(n) = f_{231}(n) = \binom{n}{3}$
$S_n(132, 312)$	$f_{123}(n) = f_{213}(n) = f_{231}(n) = f_{321}(n) = \frac{2^n}{8}\binom{n}{3}$		$f_{321}(n) = (n-2)\binom{n}{3}$
$S_n(132, 321)$	$f_{213}(n) = f_{231}(n) = f_{312}(n) = \binom{n+2}{5}$		$f_{132}(n) = f_{213}(n) = \binom{n+1}{4}$
	$f_{123}(n) = \frac{7n^5}{120} - \frac{n^4}{3} + \frac{17n^3}{24} - \frac{2n^2}{3} + \frac{7}{30}$	$S_n(123, 231, 312)$	$f_{321}(n) = \frac{1}{12}n(n-2)(n-1)^2$

Table 1: The pattern count on doubly and triply restricted permutations.

and showed both algebraically and bijectively that

$$f_{231}(S_n(132)) = f_{312}(S_n(132)) = f_{213}(S_n(132)).$$

Rudolph [7] proved equipopularity relations between general length k patterns on 132-avoiding permutations based on the structure of their corresponding binary plane trees. Moreover, Homberger [6] also presented exact formulae for each length 3 pattern in the set $S_n(123)$. Therefore, the singly restricted permutations have been well studied by previous work, whereas it remains open for multiply restricted permutations, e.g., $S_n(123, 132)$.

In this paper, we are interested in pattern count on multiply restricted permutations $S_n(R)$ for $R \subset S_3$, especially for double and triple restrictions. We derive explicit formulae or generating functions for the occurrences of each length 3 pattern in multiply restricted permutations, and the detailed results are summarized in Table 1. Also, we present some combinatorial interpretations for non-trivial pattern relationships. It is trivial to consider the restricted permutations of higher multiplicity since there are only finite permutations, as shown in [8]. Therefore, this work presents a complete study on length 3 patterns of multiply restricted permutations.

2 Doubly Restricted Permutations

For $\sigma \in S_n$, the following three operations are very useful in pattern avoiding enumeration. The complement of σ is given by $\sigma^c = (n+1-\sigma_1)(n+1-\sigma_2) \cdots (n+1-\sigma_n)$, its reverse is defined as $\sigma^r = \sigma_n \cdots \sigma_2 \sigma_1$ and the inverse σ^{-1} is the group theoretic inverse permutation. For any set of permutations R , let R^c be the set obtained by complementing each element of R , and the set R^r and R^{-1} are defined analogously.

Lemma 2.1. *Let $R \subseteq S_k$ be any set of permutations in S_k , and $\sigma \in S_n$, we have*

$$\sigma \in S_n(R) \Leftrightarrow \sigma^c \in S_n(R^c) \Leftrightarrow \sigma^r \in S_n(R^r) \Leftrightarrow \sigma^{-1} \in S_n(R^{-1}).$$

From Lemma 2.1 and the known results on $S_n(123)$ and $S_n(132)$, we can obtain each length 3 pattern count in the singly restricted permutations $S_n(r)$ for $r = 213, 231, 312$ and 321 . In this section, we focus on the number of length 3 patterns in all doubly restricted permutations.

A composition of n is an expression of n as an ordered sum of positive integers, and we say that c has k parts or c is a k -composition if there are exactly k summands appeared in a composition c . Denote by $\mathcal{C}_n$ and $\mathcal{C}_{n,k}$ the set of all compositions of n and the set of k -compositions of n , respectively. It is known that $|\mathcal{C}_n| = 2^{n-1}$ and $|\mathcal{C}_{n,k}| = \binom{n-1}{k-1}$ for $n \geq 1$ with $|\mathcal{C}_0| = 1$. For more details, see [9].

We begin with some enumerative results on compositions.

Lemma 2.2. *For $n \geq 1$, we have*

$$a(n) := \sum_{\substack{k \geq 1 \\ c_1 + \dots + c_{k-1} + c_k = n}} c_k = 2^n - 1, \quad (2.1)$$

$$b(n) := \sum_{\substack{k \geq 1 \\ c_1 + \dots + c_{k-1} + c_k = n}} c_k(c_k - 1) = 2^{n+1} - 2n - 2, \quad (2.2)$$

where the sum takes over all compositions of n .

Proof. For $c_k = m$, we can regard $c_1 + c_2 + \dots + c_{k-1}$ as a composition of $n - m$. It is easy to get that the number of compositions of $n - m$ is 2^{n-m-1} for $1 \leq m \leq n - 1$ and the number of empty compositions is one. This follows that

$$\begin{aligned} a(n) &= n + \sum_{m=1}^{n-1} m 2^{n-m-1}, \\ b(n) &= n(n-1) + \sum_{m=1}^{n-1} m(m-1) 2^{n-m-1}. \end{aligned}$$

Let $g(x) = \sum_{i=0}^{n-1} x^i = \frac{1-x^n}{1-x}$, and we have

$$\begin{aligned} g'(x) &= \sum_{i=1}^{n-1} i x^{i-1} = \frac{(n-1)x^n - nx^{n-1} + 1}{(1-x)^2}, \\ g''(x) &= \sum_{i=1}^{n-1} i(i-1)x^{i-2} = \frac{(3n-n^2-2)x^n + (2n^2-4n)x^{n-1} + (n-n^2)x^{n-2} + 2}{(1-x)^3}. \end{aligned}$$

By setting $x = 1/2$ in $g'(x)$ and $g''(x)$, we get

$$\begin{aligned} g'(1/2) &= 2^2((n-1)2^{-n} - n2^{-n+1} + 1), \\ g''(1/2) &= 2^3[(3n - n^2 - 2)2^{-n} + (2n^2 - 4n)2^{-n+1} + (n - n^2)2^{-n+2} + 2]. \end{aligned}$$

Observing $a(n) = 2^{n-2}g'(1/2) + n$ and $b(n) = 2^{n-3}g''(1/2) + n(n-1)$, this lemma follows as desired. ■

Lemma 2.3. For $n \geq 1$, we have

$$c(n) := \sum_{\substack{k \geq 1 \\ c_1 + \dots + c_{k-1} + c_k = n}} k = (n+1)2^{n-2}, \quad (2.3)$$

$$d(n) := \sum_{\substack{k \geq 1 \\ c_1 + \dots + c_{k-1} + c_k = n}} k(k-1) = (n^2 + n - 2)2^{n-3}, \quad (2.4)$$

where the sum takes over all compositions of n .

Proof. Since the number of compositions of n with k parts is $\binom{n-1}{k-1}$, we have

$$c(n) = \sum_{k=1}^n k \binom{n-1}{k-1} \text{ and } d(n) = \sum_{k=1}^n k(k-1) \binom{n-1}{k-1}.$$

Let $h(x) = x \sum_{i=1}^n \binom{n-1}{i-1} x^{i-1} = x(1+x)^{n-1}$. Then

$$\begin{aligned} h'(x) &= \sum_{i=1}^n i \binom{n-1}{i-1} x^{i-1} = (nx+1)(1+x)^{n-2}, \\ h''(x) &= \sum_{i=1}^n i(i-1) \binom{n-1}{i-1} x^{i-2} = (n^2x + n(2-x) - 2)(1+x)^{n-3}. \end{aligned}$$

We complete the proof by putting $x = 1$ in the above formulae. ■

Based on Lemma 2.1, Simion and Schmidt [8] showed that the pairs of patterns among the total $\binom{6}{2} = 15$ cases fall into the following 6 classes.

Proposition 2.4 ([8]). For every symmetric group S_n ,

1. $|S_n(123, 132)| = |S_n(123, 213)| = |S_n(231, 321)| = |S_n(312, 321)| = 2^{n-1};$
2. $|S_n(132, 213)| = |S_n(231, 312)| = 2^{n-1};$
3. $|S_n(132, 231)| = |S_n(213, 312)| = 2^{n-1};$
4. $|S_n(132, 312)| = |S_n(213, 231)| = 2^{n-1};$
5. $|S_n(132, 321)| = |S_n(123, 231)| = |S_n(123, 312)| = |S_n(213, 321)| = \binom{n}{2} + 1;$
6. $|S_n(123, 321)| = 0$ for $n \geq 5$.

Therefore, it is sufficient to consider the pattern count of the first set for each class in the subsequent sections, and the other sets can be obtained by taking the complement or reverse or inverse of the known results.

2.1 Pattern Count on (123, 132)-Avoiding Permutations

We first present a bijection between $S_n(123, 132)$ and $\mathcal{C}_n$ as follows:

Lemma 2.5. *There is a bijection φ_1 between the sets $S_n(123, 132)$ and $\mathcal{C}_n$.*

Proof. For any given $\sigma \in S_n(123, 132)$, let $\sigma_{i_1}, \sigma_{i_2}, \dots, \sigma_{i_k}$ be the k right to left maxima with $i_1 < i_2 < \dots < i_k$, which yields that $c = i_1 + (i_2 - i_1) + \dots + (i_{k-1} - i_{k-2}) + (i_k - i_{k-1})$ is a composition of n since $i_k = n$. On the converse, let $m_i = n - (c_1 + \dots + c_{i-1})$ for any given $n = c_1 + c_2 + \dots + c_k \in \mathcal{C}_n$, and we set $\tau_i = m_i - 1, m_i - 2, \dots, m_i - c_i + 1, m_i$ for $1 \leq i \leq k$. Therefore, $\sigma = \tau_1, \tau_2, \dots, \tau_k \in S_n(123, 132)$ is as desired. For example, we have $\sigma = 897543612$ for the composition $9 = 2 + 1 + 4 + 2$. ■

For a pattern q , we denote by $f_q(n) := \sum_{\sigma \in S_n(123, 132)} f_q(\sigma)$, i.e., the number of occurrences of the pattern q in $S_n(123, 132)$. For simplicity, we will use this notation in subsequent sections when the set in question is unambiguous. For convenience, we denote by $\tau_i > \tau_j$ if all the elements in the subsequence τ_i are larger than all the elements in subsequence τ_j . Based on Lemma 2.5, we have

Proposition 2.6. *For $n \geq 3$,*

$$f_{213}(n) = \sum_{\substack{k \geq 1 \\ c_1 + c_2 + \dots + c_k = n}} \sum_{i=1}^k \binom{c_i - 1}{2}, \quad (2.5)$$

$$f_{231}(n) = \sum_{\substack{k \geq 1 \\ c_1 + c_2 + \dots + c_k = n}} \sum_{i=1}^{k-1} \sum_{j=i+1}^k c_j(c_i - 1). \quad (2.6)$$

Proof. For each permutation $\sigma \in S_n(123, 132)$ with $\varphi_1(\sigma) = c_1 + c_2 + \dots + c_k$, we can rewrite σ as $\sigma = \tau_1, \tau_2, \dots, \tau_k$ from Lemma 2.5. For $j > i$, since $\tau_i > \tau_j$, and the elements except the last one are decreasing in τ_i , the pattern 213 can only occur in every subsequence τ_i . Thus, we have $\binom{c_i - 1}{2}$ choices to choose two elements in τ_i to play the role of 21, and the last element of τ_i plays the role of 3, summing up all the number of 213-patterns in subsequences $\tau_1, \tau_2, \dots, \tau_k$ gives the formula (2.5).

For the pattern 231, we have $c_i - 1$ choices in the subsequence τ_i to choose one element playing the role of 2 and one choice (always the last element of τ_i) for 3, and then, we have $c_{i+1} + \dots + c_k$ choices to choose one element for the role of 1 since all the elements after τ_i are smaller than those in τ_i . Summing up all the number of 231-patterns according to the position of 3 gives the formula (2.6). ■

Based on the previous analysis, we now present our first main results of the explicit formulae for pattern count in the set $S_n(123, 132)$.

Theorem 2.7. For $n \geq 3$, in the set $S_n(123, 132)$, we have

$$f_{213}(n) = (n-3)2^{n-2} + 1, \quad (2.7)$$

$$f_{231}(n) = f_{312}(n) = (n^2 - 5n + 8)2^{n-3} - 1, \quad (2.8)$$

$$f_{321}(n) = (n^3/3 - 2n^2 + 14n/3 - 5)2^{n-2} + 1. \quad (2.9)$$

Proof. From $S_3(123, 132) = \{213, 231, 312, 321\}$, it is obvious that

$$f_{213}(3) = f_{231}(3) = 1.$$

To prove formula (2.7), from Prop 2.6, we observe that, for $n \geq 3$

$$f_{213}(n+1) = \sum_{\substack{k \geq 1, c_k = 1 \\ c_1 + c_2 + \dots + c_k = n+1}} \sum_{i=1}^k \binom{c_i - 1}{2} + \sum_{\substack{k \geq 1, c_k \geq 2 \\ c_1 + c_2 + \dots + c_k = n+1}} \sum_{i=1}^k \binom{c_i - 1}{2}.$$

If $c_k = 1$, then $k \geq 2$, and we further have

$$\sum_{\substack{k \geq 1, c_k = 1 \\ c_1 + c_2 + \dots + c_k = n+1}} \sum_{i=1}^k \binom{c_i - 1}{2} = \sum_{\substack{k-1 \geq 1 \\ c_1 + c_2 + \dots + c_{k-1} = n}} \sum_{i=1}^{k-1} \binom{c_i - 1}{2} = f_{213}(n).$$

If $c_k \geq 2$, then we set $c_k = 1 + r_k$, and from Lemma 2.2, it holds that

$$\begin{aligned} \sum_{\substack{k \geq 1, c_k \geq 2 \\ c_1 + c_2 + \dots + c_k = n+1}} \sum_{i=1}^k \binom{c_i - 1}{2} &= \sum_{\substack{k \geq 1 \\ c_1 + \dots + c_{k-1} + r_k = n}} \left[\sum_{i=1}^{k-1} \binom{c_i - 1}{2} + \binom{r_k - 1}{2} + (r_k - 1) \right] \\ &= f_{213}(n) + \sum_{\substack{k \geq 1 \\ c_1 + \dots + c_{k-1} + r_k = n}} (r_k - 1) \\ &= f_{213}(n) + a(n) - \sum_{\substack{k \geq 1 \\ c_1 + \dots + c_{k-1} + r_k = n}} 1 = f_{213}(n) + a(n) - 2^{n-1}. \end{aligned}$$

Combining the above two cases, we have

$$f_{213}(n+1) = 2f_{213}(n) + 2^{n-1} - 1.$$

This proves formula (2.7) by solving the recurrence with initial value $f_{213}(3) = 1$.

To prove formula (2.8), first observe that from Lemma 2.1, $\sigma \in S_n(123, 132) \Leftrightarrow \sigma^{-1} \in S_n(123, 132)$, this implies the first equality of formula (2.8) directly since $231^{-1} = 312$. While for the second equality of formula (2.8), by Prop 2.6, we have

$$f_{231}(n+1) = \sum_{\substack{k \geq 1, c_k = 1 \\ c_1 + c_2 + \dots + c_k = n+1}} \sum_{i=1}^{k-1} \sum_{j=i+1}^k c_j(c_i - 1) + \sum_{\substack{k \geq 1, c_k \geq 2 \\ c_1 + c_2 + \dots + c_k = n+1}} \sum_{i=1}^{k-1} \sum_{j=i+1}^k c_j(c_i - 1).$$

If $c_k = 1$, then $k \geq 2$, and from Lemma 2.3 we have

$$\begin{aligned}
\sum_{\substack{k \geq 1, c_k = 1 \\ c_1 + c_2 + \dots + c_k = n+1}} \sum_{i=1}^{k-1} \sum_{j=i+1}^k c_j(c_i - 1) &= \sum_{\substack{k-1 \geq 1 \\ c_1 + \dots + c_{k-1} = n}} \sum_{i=1}^{k-2} \sum_{j=i+1}^{k-1} c_j(c_i - 1) + \sum_{\substack{k-1 \geq 1 \\ c_1 + \dots + c_{k-1} = n}} \sum_{i=1}^{k-1} (c_i - 1) \\
&= f_{231}(n) + \sum_{\substack{k-1 \geq 1 \\ c_1 + \dots + c_{k-1} = n}} n - (k-1) \\
&= f_{231}(n) - c(n) + n2^{n-1}.
\end{aligned}$$

If $c_k \geq 2$, then we set $c'_k = c_k - 1$ and $c'_i = c_i$ for $1 \leq i \leq k-1$. This holds that

$$\begin{aligned}
\sum_{\substack{k \geq 1, c_k \geq 2 \\ c_1 + \dots + c_k = n+1}} \sum_{i=1}^{k-1} \sum_{j=i+1}^k c_j(c_i - 1) &= \sum_{\substack{k \geq 1 \\ c'_1 + \dots + c'_k = n}} \sum_{i=1}^{k-1} \sum_{j=i+1}^k c'_j(c'_i - 1) + \sum_{\substack{k \geq 1 \\ c'_1 + \dots + c'_k = n}} \sum_{i=1}^{k-1} (c'_i - 1) \\
&= f_{231}(n) + \sum_{\substack{k \geq 1 \\ c'_1 + \dots + c'_k = n}} (n - c'_k - k + 1) \\
&= f_{231}(n) - a(n) - c(n) + (n+1)2^{n-1},
\end{aligned}$$

where the last equality holds from Lemma 2.2 and Lemma 2.3. Therefore, after simplification, we have

$$f_{231}(n+1) = 2f_{231}(n) + (n-2)2^{n-1} + 1,$$

which proves the second equality of formula (2.8) by using $f_{231}(3) = 1$.

Note that the total number of all length 3 patterns in a permutation $\sigma \in S_n$ is $\binom{n}{3}$, for the set $S_n(123, 132)$, this gives the relation

$$f_{213}(n) + 2f_{231}(n) + f_{321}(n) = \binom{n}{3} 2^{n-1}.$$

Thus formula (2.9) is a direct computation of the above equation. This completes the proof. $\blacksquare$

The first few values of $f_q(S_n(123, 132))$ for q of length 3 are shown below.

n	f_{123}	f_{132}	f_{213}	f_{231}	f_{312}	f_{321}	n	f_{123}	f_{132}	f_{213}	f_{231}	f_{312}	f_{321}
3	0	0	1	1	1	1	6	0	0	49	111	111	369
4	0	0	5	7	7	13	7	0	0	129	351	351	1409
5	0	0	17	31	31	81	8	0	0	321	1023	1023	4801

2.2 Pattern Count on (132, 213)-Avoiding Permutations

We begin with the following correspondence between (132, 213)-avoiding permutations and compositions of n .

Lemma 2.8. *There is a bijection φ_2 between the sets $S_n(132, 213)$ and $\mathcal{C}_n$.*

Proof. Given $\sigma \in S_n(132, 213)$, let $\sigma_{i_1}, \sigma_{i_2}, \dots, \sigma_{i_k}$ be the k right to left maxima with $i_1 < i_2 < \dots < i_k$. This follows that $c = i_1 + (i_2 - i_1) + \dots + (i_{k-1} - i_{k-2}) + (i_k - i_{k-1})$ is a composition of n since $i_k = n$. On the converse, given $n = c_1 + c_2 + \dots + c_k \in \mathcal{C}_n$, let $m_i = n - (c_1 + \dots + c_{i-1})$ and $\tau_i = m_i - c_i + 1, m_i - c_i + 2, \dots, m_i - 1, m_i$ for $1 \leq i \leq k$. We set $\sigma = \tau_1, \tau_2, \dots, \tau_k$, and it is easy to check that $\sigma \in S_n(132, 213)$. For example, for the composition $9 = 3 + 3 + 1 + 2$, we get $\sigma = 789456312$. ■

Based on the above lemma, we have

Proposition 2.9. *For $n \geq 3$,*

$$f_{123}(n) = \sum_{\substack{k \geq 1 \\ c_1 + c_2 + \dots + c_k = n}} \sum_{i=1}^k \binom{c_i}{3}, \quad (2.10)$$

$$f_{231}(n) = \sum_{\substack{k \geq 1 \\ c_1 + c_2 + \dots + c_k = n}} \sum_{i=1}^{k-1} \sum_{j=i+1}^k c_j \binom{c_i}{2}. \quad (2.11)$$

Proof. For a permutation $\sigma \in S_n(132, 213)$ with $\varphi_2(\sigma) = c_1 + c_2 + \dots + c_k$, we rewrite σ as $\sigma = \tau_1, \tau_2, \dots, \tau_k$ by Lemma 2.8, and we see that the pattern 123 can only occur in every subsequence τ_i since $\tau_i > \tau_j$ for $j > i$ and the elements in τ_i are increasing. Thus, we have $\binom{c_i}{3}$ choices to choose three elements in τ_i to play the role of 123, and formula (2.10) follows by summing all 123-patterns in subsequences $\tau_1, \tau_2, \dots, \tau_k$.

For pattern 231, we have $\binom{c_i}{2}$ choices in the subsequence τ_i to choose two elements to play the role of 23, after this we have $c_{i+1} + \dots + c_k$ choices to choose one element in $\tau_{i+1}, \dots, \tau_k$ for the role of 1 since $\tau_j < \tau_i$ for all $j > i$. Summing up all the number of 231-patterns according to the positions of 23 gives the formula (2.11). ■

Theorem 2.10. *For $n \geq 3$, in the set $S_n(132, 213)$, we have*

$$f_{123}(n) = (n - 4)2^{n-1} + n + 2, \quad (2.12)$$

$$f_{231}(n) = f_{312}(n) = (n^2 - 7n + 16)2^{n-2} - n - 4, \quad (2.13)$$

$$f_{321}(n) = (n^3/3 - 3n^2 + 38n/3 - 24)2^{n-2} + n + 6. \quad (2.14)$$

Proof. From Prop 2.9, we have

$$f_{123}(n+1) = \sum_{\substack{k \geq 1, c_k = 1 \\ c_1 + c_2 + \dots + c_k = n+1}} \sum_{i=1}^k \binom{c_i}{3} + \sum_{\substack{k \geq 1, c_k \geq 2 \\ c_1 + c_2 + \dots + c_k = n+1}} \sum_{i=1}^k \binom{c_i}{3}.$$

If $c_k = 1$, then $k \geq 2$, and we have

$$\sum_{\substack{k \geq 1, c_k = 1 \\ c_1 + c_2 + \dots + c_k = n+1}} \sum_{i=1}^k \binom{c_i}{3} = \sum_{\substack{k-1 \geq 1 \\ c_1 + c_2 + \dots + c_{k-1} = n}} \sum_{i=1}^{k-1} \binom{c_i}{3} = f_{123}(n).$$

If $c_k \geq 2$, then we set $c_k = 1 + r_k$, where $r_k \geq 1$, and this follows

$$\begin{aligned} \sum_{\substack{k \geq 1, c_k \geq 2 \\ c_1 + c_2 + \dots + c_k = n+1}} \sum_{i=1}^k \binom{c_i}{3} &= \sum_{\substack{k \geq 1 \\ c_1 + c_2 + \dots + c_{k-1} + r_k = n}} \left[\sum_{i=1}^{k-1} \binom{c_i}{3} + \binom{r_k}{3} + \frac{r_k(r_k - 1)}{2} \right] \\ &= f_{123}(n) + b(n)/2. \end{aligned}$$

Combining the above two cases, we get that

$$f_{123}(n+1) = 2f_{123}(n) + 2^n - n - 1,$$

and formula (2.12) holds by solving the recurrence with initial value $f_{123}(3) = 1$.

From Lemma 2.1, we see that $\sigma \in S_n(132, 213) \Leftrightarrow \sigma^{-1} \in S_n(132, 213)$, and this follows that $f_{231}(n) = f_{312}(n)$ from $231^{-1} = 312$.

To calculate $f_{231}(n)$, we have by using Prop 2.9 again

$$f_{231}(n+1) = \sum_{\substack{k \geq 1, c_k = 1 \\ c_1 + c_2 + \dots + c_k = n+1}} \sum_{i=1}^{k-1} \sum_{j=i+1}^k c_j \binom{c_i}{2} + \sum_{\substack{k \geq 1, c_k \geq 2 \\ c_1 + c_2 + \dots + c_k = n+1}} \sum_{i=1}^{k-1} \sum_{j=i+1}^k c_j \binom{c_i}{2}.$$

If $c_k = 1$, then $k \geq 2$, and

$$\begin{aligned} \sum_{\substack{k \geq 1, c_k = 1 \\ c_1 + c_2 + \dots + c_k = n+1}} \sum_{i=1}^{k-1} \sum_{j=i+1}^k c_j \binom{c_i}{2} &= \sum_{\substack{k-1 \geq 1 \\ c_1 + c_2 + \dots + c_{k-1} = n}} \sum_{i=1}^{k-1} \binom{c_i}{2} \left[\sum_{j=i+1}^{k-1} c_j + 1 \right] \\ &= f_{231}(n) + \alpha(n), \end{aligned}$$

where $\alpha(n) = \sum_{\substack{k \geq 1 \\ c_1 + c_2 + \dots + c_k = n}} \sum_{i=1}^k \binom{c_i}{2}$. We further have

$$\begin{aligned} \alpha(n) &= \sum_{\substack{k \geq 1 \\ c_1 + \dots + c_k = n}} \sum_{i=1}^k \left[\binom{c_i - 1}{2} + c_i - 1 \right] = f_{213}(S_n(123, 132)) + \sum_{\substack{k \geq 1 \\ c_1 + \dots + c_k = n}} (n - k) \\ &= f_{213}(S_n(123, 132)) - c(n) + n2^{n-1}, \end{aligned}$$

in which we use the derived formula $f_{213}(S_n(123, 132)) = \sum_{\substack{k \geq 1 \\ c_1 + \dots + c_k = n}} \sum_{i=1}^k \binom{c_i - 1}{2}$.

If $c_k \geq 2$, then we set $c'_k = c_k - 1$ and $c'_i = c_i$ for $1 \leq i \leq k-1$. This holds that

$$\begin{aligned} \sum_{\substack{k \geq 1, c_k \geq 2 \\ c_1 + \dots + c_k = n+1}} \sum_{i=1}^{k-1} \sum_{j=i+1}^k c_j \binom{c_i}{2} &= \sum_{\substack{k \geq 1 \\ c'_1 + \dots + c'_k = n}} \sum_{i=1}^{k-1} \sum_{j=i+1}^k c'_j \binom{c'_i}{2} + \sum_{\substack{k \geq 1 \\ c'_1 + \dots + c'_k = n}} \sum_{i=1}^{k-1} \binom{c'_i}{2} \\ &= f_{231}(n) + \beta(n), \end{aligned}$$

where $\beta(n) = \sum_{\substack{k \geq 1 \\ c_1 + \dots + c_k = n}} \sum_{i=1}^{k-1} \binom{c_i}{2}$. Further, we can rewrite $\beta(n)$ as

$$\begin{aligned} \beta(n) &= \sum_{\substack{k \geq 1 \\ c_1 + \dots + c_k = n}} \sum_{i=1}^k \binom{c_i}{2} - \sum_{\substack{k \geq 1 \\ c_1 + \dots + c_k = n}} \frac{c_k(c_k - 1)}{2} \\ &= \alpha(n) - b(n)/2 = f_{213}(S_n(123, 132)) - c(n) - b(n)/2 + n2^{n-1}. \end{aligned}$$

Substituting the known formulae for $f_{213}(S_n(123, 132))$, $c(n)$ and $b(n)$, we get that

$$f_{231}(n+1) = 2f_{231}(n) + (2n-6)2^{n-1} + n+3,$$

and formula (2.14) holds by solving this recurrence with initial condition $f_{213}(3) = 1$.

Finally, formula (2.14) follows from $f_{123}(n) + 2f_{231}(n) + f_{321}(n) = \binom{n}{3}2^{n-1}$, and this completes the proof. $\blacksquare$

The first few values of $f_q(S_n(132, 213))$ for q of length 3 are shown below.

n	f_{123}	f_{132}	f_{213}	f_{231}	f_{312}	f_{321}	n	f_{123}	f_{132}	f_{213}	f_{231}	f_{312}	f_{321}
3	1	0	0	1	1	1	6	72	0	0	150	150	268
4	6	0	0	8	8	10	7	201	0	0	501	501	1037
5	23	0	0	39	39	59	8	522	0	0	1524	1524	3598

2.3 Pattern Count on (132, 231)-Avoiding Permutations

Theorem 2.11. *For $n \geq 3$, in the set $S_n(132, 231)$, we have*

$$f_{123}(n) = f_{213}(n) = f_{312}(n) = f_{321}(n) = \binom{n}{3}2^{n-3}. \quad (2.15)$$

Proof. For each $\sigma \in S_n(132, 231)$, we observe that n must lie in the beginning or the end of σ , and $n-1$ must lie in the beginning or the end of $\sigma \setminus \{n\}$, ..., and so on. Here $\sigma \setminus \{n\}$ denotes the sequence obtained from σ by deleting element n . Based on such observation, suppose abc is a length 3 pattern in $S_n(132, 231)$, and set $[n] \setminus \{a, b, c\} := \{r_1 > r_2 > \dots > r_{n-4} > r_{n-3}\}$. We can construct a permutation in the set $S_n(132, 231)$ which contains an abc pattern as follows:

Start with the subsequence $\sigma^0 := abc$, and for i from 1 to $n-3$, σ^i is obtained from σ^{i-1} by inserting r_i into it.

- if there are at least two elements in σ^{i-1} smaller than r_i , then choose the two elements A and B such that A is the leftmost one and B is the rightmost one. We put r_i immediately to the left of A or immediately to the right of B ;

- if there is only one element A in σ^{i-1} such that $A < r_i$, then we can put r_i immediately to the left or to the right of A ;
- if all the elements in σ^{i-1} are larger than r_i , then choose A the least one, and put r_i immediately to the left or to the right of A .

Finally, we set $\sigma := \sigma^{n-3}$ and $\sigma \in S_n(132, 231)$ from the above construction. Moreover, there are $\binom{n}{3}$ choices to choose abc , and the number of permutations having abc as a pattern is 2^{n-3} since each r_i has 2 choices. This completes the proof. $\blacksquare$

Here we give an illustration of constructing a permutation in $S_8(132, 231)$ which contains the pattern $abc = 256$. Set $\sigma^0 := 256$, we may have $\sigma^1 = 8256$, $\sigma^2 = 87256$, $\sigma^3 = 872456$, $\sigma^4 = 8732456$, $\sigma := \sigma^5 = 87321456$.

We could also provide combinatorial proofs for the phenomenon $f_{123}(n) = f_{213}(n) = f_{312}(n) = f_{321}(n)$. From Lemma 2.1, we have $\sigma \in S_n(132, 231) \Leftrightarrow \sigma^r \in S_n(132, 231)$, and this follows $f_{123}(n) = f_{321}(n)$ and $f_{213}(n) = f_{312}(n)$ from $123^r = 321$ and $213^r = 312$, respectively. It remains to give a bijection for $f_{213}(n) = f_{123}(n)$, and our following construction is motivated from Bóna [2].

A binary plane tree is a rooted unlabeled tree in which each vertex has at most two children, and each child is a left child or a right child of its parent. For each $\sigma \in S_n(132)$, we construct a binary plane tree $T(\sigma)$ as follows: the root of $T(\sigma)$ corresponds to the entry n of σ , the left subtree of the root corresponds to the string of entries of σ on the left of n , and the right subtree of the root corresponds to the string of entries of σ on the right of n . Both subtrees are constructed recursively by the same rule. For more details, see [1, 2, 7].

A left descendant (resp. right descendant) of a vertex x in a binary plane tree is a vertex in the left (resp. right) subtree of x . The left (resp. right) subtree of x does not contain x itself. Similarly, an ascendant of a vertex x in a binary plane tree is a vertex whose subtree contains x . Given a tree T and a vertex $v \in T$, let T_v be the subtree of T with v as the root. Let R be an occurrence of the pattern 123 in $\sigma \in S_n(132)$, and let R_1, R_2, R_3 be the three vertices of $T(\sigma)$ that correspond to R , going left to right. Then, R_1 is a left descendant of R_2 , and R_2 is a left descendant of R_3 .

From the above correspondence, we see that for $\sigma \in S_n(132, 231)$, $T(\sigma)$ is a binary plane tree on n vertices such that each vertex has at most one child from the forbiddance of the pattern 231. For simplicity, denote by $\mathcal{T}_n$ the set of such binary plane trees on n vertices. Let Q be an occurrence of the pattern 213 in $\sigma \in S_n(132, 231)$, and let Q_2, Q_1, Q_3 be the three vertices of $T(\sigma)$ that correspond to Q , going left to right. From the characterization of trees in $\mathcal{T}_n$, Q_2 is a left descendant of Q_3 , and Q_1 is a right descendant of Q_2 .

Let $\mathcal{A}_n$ be the set of binary plane trees in $\mathcal{T}_n$ where three vertices forming a 213-pattern are colored black. Let $\mathcal{B}_n$ be the set of all binary plane trees in $\mathcal{T}_n$ where three vertices forming a 123-pattern are colored black. We will define a map $\rho : \mathcal{A}_n \rightarrow \mathcal{B}_n$

as follows. Given a tree $T \in \mathcal{A}_n$ with Q_2, Q_1, Q_3 being the three black vertices as a 213-pattern, define $\rho(T)$ be the tree obtained by changing the right subtree of Q_2 to be its left subtree. See Figure 1 for an illustration.




Figure 1: The bijection ρ .

In the tree $\rho(T)$, the relative positions of Q_2 and Q_3 keep the same, and Q_1 is a left descendant of Q_2 . Therefore, the three black points $Q_1Q_2Q_3$ form a 123-pattern in $\rho(T)$, and $\rho(T) \in \mathcal{B}_n$. It is easy to describe the converse and we omit here.

The first few values of $f_q(S_n(132, 231))$ for q of length 3 are shown below.

n	f_{123}	f_{132}	f_{213}	f_{231}	f_{312}	f_{321}	n	f_{123}	f_{132}	f_{213}	f_{231}	f_{312}	f_{321}
3	1	0	1	0	1	1	6	160	0	160	0	160	160
4	8	0	8	0	8	8	7	560	0	560	0	560	560
5	40	0	40	0	40	40	8	1792	0	1792	0	1792	1792

2.4 Pattern Count on (132, 312)-Avoiding Permutations

We begin with a correspondence between (132, 312)-avoiding permutations and compositions of n .

Theorem 2.12. *There is a bijection φ_4 between the sets $S_n(132, 312)$ and $\mathcal{C}_n$.*

Proof. For $\sigma \in S_n(132, 312)$, let $\sigma_{i_1}, \sigma_{i_2}, \dots, \sigma_{i_k}$ be the k left to right maxima with $i_1 < i_2 < \dots < i_k$, and thus, $c = (i_2 - i_1) + (i_3 - i_2) + \dots + (i_k - i_{k-1}) + (n + 1 - i_k)$ is a composition of n from $i_1 = 1$. On the converse, let $n = c_k + c_{k-1} + \dots + c_2 + c_1 \in \mathcal{C}_n$. For $1 \leq i \leq k$, if $c_i = 1$ then set $\tau_i = n - i + 1$; otherwise, set $m_i = c_1 + \dots + c_{i-1} - i + 2$ and $\tau_i = n - i + 1, m_i + c_i - 2, \dots, m_i + 1, m_i$. It is easy to get $\sigma = \tau_k, \tau_{k-1}, \dots, \tau_2, \tau_1 \in S_n(132, 312)$ as desired. For example, if $9 = 3 + 1 + 2 + 3$, then $\sigma = 654783921$. ■

Proposition 2.13. *For $n \geq 3$, we have*

$$f_{123}(n) = \sum_{\substack{k \geq 1 \\ c_1 + c_2 + \dots + c_k = n}} \sum_{i=1}^{k-2} c_i \binom{k-i}{2}. \quad (2.16)$$

Proof. Given a permutation $\sigma = \tau_k, \dots, \tau_2, \tau_1$ in $S_n(132, 312)$ whose composition is given by $n = c_k + c_{k-1} + \dots + c_2 + c_1$, we see that the first element in τ_i is larger than all the elements in τ_j , whereas the other elements in τ_i are smaller than the elements in τ_j for $i + 1 \leq j \leq k$. The left to right maxima form an increasing subsequence and the other elements form a decreasing subsequence. Thus we have c_i choices to choose one element in τ_{k-i+1} to play the role of 1, and then $\binom{k-i}{2}$ choices to choose two left to right maxima in τ_j for $j < k - i + 1$ to play the role of 23, summing up all the number of 123 patterns in subsequences $\tau_k, \dots, \tau_2, \tau_1$ gives the formula (2.16). $\blacksquare$

Theorem 2.14. *For $n \geq 3$, in the set $S_n(132, 312)$, we have*

$$f_{123}(n) = f_{321}(n) = \binom{n}{3} 2^{n-3}, \quad (2.17)$$

$$f_{213}(n) = f_{231}(n) = \binom{n}{3} 2^{n-3}. \quad (2.18)$$

Proof. From Lemma 2.1, we see that $\sigma \in S_n(132, 312) \Leftrightarrow \sigma^c \in S_n(132, 312)$, which follows $f_{123}(n) = f_{321}(n)$ and $f_{213}(n) = f_{231}(n)$ from $123^c = 321$ and $213^c = 231$, respectively.

To calculate $f_{123}(n)$, we have by Prop 2.13

$$f_{123}(n+1) = \sum_{\substack{k \geq 1, c_1 = 1 \\ c_1 + c_2 + \dots + c_k = n+1}} \sum_{i=1}^{k-2} c_i \binom{k-i}{2} + \sum_{\substack{k \geq 1, c_1 \geq 2 \\ c_1 + c_2 + \dots + c_k = n+1}} \sum_{i=1}^{k-2} c_i \binom{k-i}{2}.$$

If $c_1 = 1$, then $k \geq 2$, and

$$\begin{aligned} \sum_{\substack{k \geq 1, c_1 = 1 \\ c_1 + c_2 + \dots + c_k = n+1}} \sum_{i=1}^{k-2} c_i \binom{k-i}{2} &= \sum_{\substack{k-1 \geq 1 \\ c_2 + \dots + c_k = n}} \sum_{i=2}^{k-2} c_i \binom{k-i}{2} + \sum_{\substack{k-1 \geq 1 \\ c_2 + \dots + c_k = n}} \binom{k-1}{2} \\ &= f_{123}(n) + d(n)/2. \end{aligned}$$

If $c_1 \geq 2$, let $c'_1 = c_1 - 1$, $c'_i = c_i$ for $2 \leq i \leq k$, then $c'_1 \geq 1$, and

$$\begin{aligned} \sum_{\substack{k \geq 1, c_1 \geq 2 \\ c_1 + c_2 + \dots + c_k = n+1}} \sum_{i=1}^{k-2} c_i \binom{k-i}{2} &= \sum_{\substack{k \geq 1 \\ c'_1 + c'_2 + \dots + c'_k = n}} \sum_{i=1}^{k-2} c'_i \binom{k-i}{2} + \sum_{\substack{k \geq 1 \\ c'_1 + c'_2 + \dots + c'_k = n}} \binom{k-1}{2} \\ &= f_{123}(n) + \sum_{\substack{k \geq 1 \\ c'_1 + c'_2 + \dots + c'_k = n}} \left[\binom{k}{2} + 1 - k \right] \\ &= f_{123}(n) + d(n)/2 + 2^{n-1} - c(n). \end{aligned}$$

Combining the above two cases, we get that

$$f_{123}(n+1) = 2f_{123}(n) + (n^2 - n)2^{n-3},$$

and the formula (2.17) is derived by solving the recurrence with initial value $f_{123}(3) = 1$. Formula (2.18) is a direct computation of the equality $2f_{123}(n) + 2f_{213}(n) = \binom{n}{3} 2^{n-1}$. $\blacksquare$

In the subsequent section, we could also give a combinatorial interpretation for $f_{231}(n) = f_{123}(n)$ by using binary plane trees.

For $\sigma \in S_n(132, 312)$, as in previous section, we could construct a binary plane tree $T(\sigma)$ on n vertices such that each vertex that is a right descendant of some vertex does not have a left descendant from the forbiddance of the pattern 312. Denote by $\mathcal{T}_n$ the set of such trees on n vertices. Let Q be an occurrence of the pattern 231 in $\sigma \in S_n(132, 312)$, and let Q_2, Q_3, Q_1 be the three vertices of $T(\sigma)$ that correspond to Q , going left to right. Then, Q_2 is a left descendant of Q_3 , and there exists a lowest ascendant x of Q_3 or $x = Q_3$ so that Q_1 is a right descendant of x . Let $\mathcal{A}_n$ be the set of binary plane trees in $\mathcal{T}_n$ in which three vertices forming a 231-pattern are colored black. Let $\mathcal{B}_n$ be the set of all binary plane trees in $\mathcal{T}_n$ in which three vertices forming a 123-pattern are colored black. It remains to construct a map $\varrho : \mathcal{A}_n \rightarrow \mathcal{B}_n$.

Given a tree $T \in \mathcal{A}_n$ with Q_2, Q_3, Q_1 being the three black vertices as a 231-pattern, denoted by y the vertex that is the parent of x if it exists. We can see that x is the left child of y from $T \in \mathcal{A}_n$. Let $T^u := T - T_x$ be the tree obtained from T by deleting the subtree T_x , and $T^d := T_x - T_{Q_1}$ be the tree obtained from T_x by deleting the subtree T_{Q_1} . Now we can define $\varrho(T)$ be the tree obtained from T by first interchanging Q_1 as the left child of y , then adjoining the subtree T^d as the left subtree of the vertex Q_1 and keeping all three black vertices the same if y exists. Otherwise, we adjoin the subtree T^d as the left subtree of the vertex Q_1 directly. An illustration is given in Figure 2.




Figure 2: The bijection ϱ .

In the tree $\varrho(T)$, the relative positions of Q_2 and Q_3 are unchanged, and Q_3 is a left descendant of Q_1 , thus the three black points $Q_2Q_3Q_1$ form a 123-pattern in $\varrho(T)$, and $\varrho(T) \in \mathcal{B}_n$. It is easy to describe the converse map and we omit here.

The first few values of $f_q(S_n(132, 312))$ for q of length 3 are shown below.

n	f_{123}	f_{132}	f_{213}	f_{231}	f_{312}	f_{321}	n	f_{123}	f_{132}	f_{213}	f_{231}	f_{312}	f_{321}
3	1	0	1	1	0	1	6	160	0	160	160	0	160
4	8	0	8	8	0	8	7	560	0	560	560	0	560
5	40	0	40	40	0	40	8	1792	0	1792	1792	0	1792

2.5 Pattern Count on (132, 321)-Avoiding Permutations

We begin with a correspondence from Simion and Schmidt [8] as follows:

Lemma 2.15. *There is a bijection φ_5 between the set $S_n(132, 321) \setminus \{\text{identity}\}$ and the set of 2-element subsets of $[n]$.*

Proof. For a permutation $\sigma \in S_n(132, 321) \setminus \{\text{identity}\}$, suppose that $\sigma_k = m$ ($k < m$), and then we define $\varphi_5(\sigma) = \{k, m\}$. On the converse, given two elements $1 \leq k < m \leq n$, set $\tau_1 = m - k + 1, m - k + 2, \dots, m - 1, m$, $\tau_2 = 1, 2, \dots, m - k$ and $\tau_3 = m + 1, m + 2, \dots, n - 1, n$. Then define $\sigma = \varphi_5^{-1}(k, m) = \tau_1, \tau_2, \tau_3$. For example, if $k = 4, m = 6$, then $\sigma = 34561278$. ■

From the above lemma, we have

Proposition 2.16. *For $n \geq 3$,*

$$f_{213}(n) = \sum_{1 \leq k < m \leq n} k(m - k)(n - m), \quad (2.19)$$

$$f_{312}(n) = \sum_{1 \leq k < m \leq n} k \binom{m - k}{2}. \quad (2.20)$$

Proof. Given a permutation $\sigma = \tau_1, \tau_2, \tau_3$ with $\varphi_5(\sigma) = \{k, m\}$ in the set $S_n(132, 321)$, we see that the elements in τ_1 , τ_2 and τ_3 are increasing, and $\tau_2 < \tau_1 < \tau_3$. We have k choices to select one element in τ_1 to play the role of 2, and then have $m - k$ choices to choose one element in τ_2 to play the role of 1, and $n - m$ choices to choose one element in τ_3 to play the role of 3. Summing up all possible k and m gives the formula (2.19).

For the pattern 312, we have k choices in the subsequence τ_1 to choose one element to play the role of 3, after this we have $\binom{m - k}{2}$ choices in the subsequence τ_2 to choose two elements to play the role of 23. Summing up all k and m gives the formula (2.20). ■

Next, we exhibit some useful formulae for calculating $f_{213}(n)$ and $f_{312}(n)$ as follows:

Lemma 2.17. *For $n \geq 2$,*

$$\begin{aligned} \sum_{k=1}^{n-1} k &= \frac{n(n-1)}{2}; & \sum_{k=1}^{n-1} k^2 &= \frac{n(n-1)(2n-1)}{6}; \\ \sum_{k=1}^{n-1} k^3 &= \frac{n^2(n-1)^2}{4}; & \sum_{k=1}^{n-1} k^4 &= \frac{n(n-1)(2n-1)(3n^2-3n+1)}{30}. \end{aligned}$$

Based on previous analysis, we are ready to give the exact formulae for the pattern count in $S_n(132, 321)$.

Theorem 2.18. For $n \geq 3$, in the set $S_n(132, 321)$, we have

$$f_{213}(n) = f_{231}(n) = f_{312}(n) = \binom{n+2}{5}, \quad (2.21)$$

$$f_{123}(n) = n(7n^4 - 40n^3 + 85n^2 - 80n + 28)/120. \quad (2.22)$$

Proof. From Lemma 2.1, we see that $\sigma \in S_n(132, 321) \Leftrightarrow \sigma^{-1} \in S_n(132, 321)$, it implies that $f_{312}(n) = f_{231}(n)$ since $312^{-1} = 231$. By using Prop 2.16, we have

$$\begin{aligned} f_{312}(n) &= \sum_{1 \leq k < m \leq n} k \binom{m-k}{2} \\ &= \sum_{k=1}^{n-1} k \sum_{m=k+1}^n \binom{m-k}{2} = \sum_{k=1}^{n-1} k \binom{n-k+1}{3} \\ &= \sum_{k=1}^{n-1} [(n^3 - n)k + (1 - 3n^2)k^2 + 2nk^3 - k^4] = \binom{n+2}{5}, \end{aligned}$$

where the last equality holds from Lemma 2.17. By using Prop 2.16 again, we have

$$\begin{aligned} f_{213}(n) &= \sum_{1 \leq k < m \leq n} k(m-k)(n-m) = \sum_{k=1}^{n-1} \sum_{m=k+1}^n k(m-k)(n-m) \\ &= \sum_{k=1}^{n-1} \sum_{m'=1}^{n-k} km'(n-m'-k) = \sum_{k=1}^{n-1} k(n-k) \sum_{m'=1}^{n-k} m' - \sum_{k=1}^{n-1} k \sum_{m'=1}^{n-k} m'^2 \\ &= \sum_{k=1}^{n-1} \left[\left(\frac{n^3}{6} - \frac{n}{6} \right) k + \left(\frac{1}{6} - \frac{n^2}{2} \right) k^2 + \frac{n}{2} k^3 - \frac{1}{6} k^4 \right] = \binom{n+2}{5}, \end{aligned}$$

where the last equality holds by simple calculation from Lemma 2.17. We complete the proof by employing the relation $2f_{231}(n) + f_{213}(n) + f_{123}(n) = \binom{n}{3} \left[\binom{n}{2} + 1 \right]$. ■

We also notice that the equality $f_{213}(n) = f_{231}(n)$ can be proved by using Bóna's bijection [2] directly on the set of binary plane trees on n vertices such that the vertex which is a right descendant of some node has no right descendants.

The first few values of $f_q(S_n(132, 321))$ for q of length 3 are shown below.

n	f_{123}	f_{132}	f_{213}	f_{231}	f_{312}	f_{321}	n	f_{123}	f_{132}	f_{213}	f_{231}	f_{312}	f_{321}
3	1	0	1	1	1	0	6	152	0	56	56	56	0
4	10	0	6	6	6	0	7	392	0	126	126	126	0
5	47	0	21	21	21	0	8	868	0	252	252	252	0

3 Triply Restricted Permutations

In this section, we study the pattern count in the simultaneous avoidance of any three patterns of length 3. Based on Lemma 2.1, Simion and Schmidt [8] showed that the pairs of patterns among the total $\binom{6}{3} = 20$ cases fall into the following 6 classes:

Proposition 3.1. *For every symmetric group S_n ,*

- (1) $|S_n(123, 132, 213)| = |S_n(231, 312, 321)| = F_{n+1}$;
- (2) $|S_n(123, 132, 231)| = |S_n(123, 213, 312)| = |S_n(132, 231, 321)| = |S_n(213, 312, 321)| = n$;
- (3) $|S_n(132, 213, 231)| = |S_n(132, 213, 312)| = |S_n(132, 231, 312)| = |S_n(213, 231, 312)| = n$;
- (4) $|S_n(123, 132, 312)| = |S_n(123, 213, 231)| = |S_n(132, 312, 321)| = |S_n(213, 231, 321)| = n$;
- (5) $|S_n(123, 231, 312)| = |S_n(132, 213, 321)| = n$;
- (6) $|S_n(R)| = 0$ for all $R \supset \{123, 321\}$ if $n \geq 5$,

where F_n is the Fibonacci number given by $F_0 = 0, F_1 = 1$ and $F_n = F_{n-1} + F_{n-2}$ for $n \geq 2$.

3.1 Pattern Count on (123, 132, 213)-Avoiding Permutations

It is known that F_{n+1} counts the number of 0-1 sequences of length $n - 1$ in which there are no consecutive ones, see [4], and we call such a sequence a Fibonacci binary word for convenience. Let B_n denote the set of all Fibonacci binary words of length n . Simion and Schmidt [8] showed that

Lemma 3.2 ([8]). *There is a bijection ψ_1 between $S_n(123, 132, 213)$ and B_{n-1} .*

Proof. Let $w = w_1w_2 \cdots w_{n-1} \in B_{n-1}$, and the corresponding permutation σ is determined as follows: For $1 \leq i \leq n - 1$, let $X_i = [n] - \{\sigma_1, \dots, \sigma_{i-1}\}$, and then set

$$\sigma_i = \begin{cases} \text{largest element in } X_i, & \text{if } s_i = 0, \\ \text{second largest element in } X_i, & \text{if } s_i = 1. \end{cases}$$

Finally, σ_n is the unique element in X_n . For example, if $w = 01001010$, then $\psi_1(w) = 978645231$. ■

Given a word $w = w_1w_2 \cdots w_n \in B_n$, we call i ($1 \leq i < n$) an ascent of w if $w_i < w_{i+1}$, and denote by $\text{asc}(w) = \{i | w_i < w_{i+1}\}$ and $\text{maj}(w) = \sum_{i \in \text{asc}(w)} i$.

Proposition 3.3. *The total number of occurrences of the pattern 312 in $S_n(123, 132, 213)$ is given by*

$$f_{312}(n) = \sum_{w \in B_{n-1}} \text{maj}(w). \quad (3.2)$$

Proof. Suppose $\sigma \in S_n(123, 132, 213)$ and $\psi_1(\sigma) = w_1w_2 \cdots w_{n-1}$. If k is an ascent of w , then $w_kw_{k+1} = 01$. We have $\sigma_k > \sigma_j$ for all $j > k$ since σ_k is the largest element

in X_k . On the other hand, there exists a unique $l > k + 1$ such that $\sigma_l > \sigma_{k+1}$ since σ_{k+1} is the second largest element in X_{k+1} . From the bijection ψ_1 , we see that for all $i \in [n - 1]$, there is at most one $j > i$ such that $\sigma_j > \sigma_i$; this implies that $\sigma_i > \sigma_{k+1}$ for all $i < k$. Thus we find that $\sigma_i \sigma_{k+1} \sigma_l$ forms a 312-pattern for all $i \leq k$, that is the ascent k will produce k 's 312-pattern in which σ_{k+1} plays the role of 1. Summing up all the ascents, we derive that there are total $\text{maj}(w)$ such patterns in σ . $\blacksquare$

Theorem 3.4. *For $n \geq 3$, in the set $S_n(123, 132, 213)$, we have*

$$\sum_{n \geq 3} f_{231}(n)x^n = \sum_{n \geq 3} f_{312}(n)x^n = \frac{x^3(1+2x)}{(1-x-x^2)^3}, \quad (3.3)$$

$$\sum_{n \geq 3} f_{321}(n)x^n = \frac{x^3(1+6x+12x^2+8x^3)}{(1-x-x^2)^4}. \quad (3.4)$$

Proof. From Lemma 2.1, we have $f_{231}(n) = f_{312}(n)$ since $\sigma \in S_n(123, 132, 213) \Leftrightarrow \sigma^{-1} \in S_n(123, 132, 213)$ and $231^{-1} = 312$. By Prop 3.3, we write

$$\sum_{n \geq 3} f_{312}(n)x^n = \sum_{n \geq 3} x^n \sum_{\sigma \in B_{n-1}} \text{maj}(w) = x \sum_{n \geq 3} \sum_{w \in B_{n-1}} \text{maj}(w)x^{n-1} = xu(x),$$

where $u(x) = \sum_{n \geq 2} \sum_{w \in B_n} \text{maj}(w)x^n$.

Let $M_n(q) = \sum_{w \in B_n} q^{\text{maj}(w)}$ and $M(x, q) = \sum_{n \geq 2} M_n(q)x^n$. Then $u(x) = \frac{\partial M(x, q)}{\partial q} \big|_{q=1}$.

Given a word $w = w_1 w_2 \cdots w_n \in B_n$, if $w_n = 0$, then $\text{maj}(w) = \text{maj}(w_1 w_2 \cdots w_{n-1})$; otherwise, $w_{n-1} w_n = 01$ and $\text{maj}(w) = \text{maj}(w_1 w_2 \cdots w_{n-2}) + (n - 1)$. Hence, we have

$$M_n(q) = M_{n-1}(q) + q^{n-1} M_{n-2}(q) \text{ for } n \geq 4, \quad (3.5)$$

with $M_2(q) = 2 + q$ and $M_3(q) = 2 + q + 2q^2$. Multiplying the recursion by x^n and summing over $n \geq 4$ yields that

$$M(x, q) - (2 + q)x^2 - (2 + q + 2q^2)x^3 = x [M(x, q) - (2 + q)x^2] + qx^2 M(xq, q).$$

Therefore

$$(1 - x)M(x, q) = qx^2 M(xq, q) + (2 + q)x^2 + 2q^2 x^3.$$

Differentiate both sides with respect to q , we get

$$(1 - x) \frac{\partial M(x, q)}{\partial q} = x^2 \left[M(xq, q) + q \frac{\partial M(xq, q)}{\partial q} \right] + x^2 + 4qx^3. \quad (3.6)$$

Setting $q = 1$ in equation (3.6) reads that

$$(1 - x)u(x) = x^2 \left[M(x, 1) + \frac{\partial M(xq, q)}{\partial q} \big|_{q=1} \right] + x^2 + 4x^3,$$

Employing the well-known generating $\sum_{n \geq 0} F_n x^n = \frac{x}{1-x-x^2}$, we have

$$M(x, 1) = \sum_{n \geq 2} |B_n| x^n = \sum_{n \geq 2} F_{n+2} x^n = \frac{x^2(3+2x)}{1-x-x^2}.$$

Further,

$$\begin{aligned} \frac{\partial M(xq, q)}{\partial q} \Big|_{q=1} &= \left(\sum_{n \geq 2} \sum_{w \in B_n} q^{n+\text{maj}(w)} x^n \right) \Big|_{q=1} \\ &= \sum_{n \geq 2} x^n \sum_{w \in B_n} (n + \text{maj}(w)) = \sum_{n \geq 2} n F_{n+2} x^n + u(x). \end{aligned}$$

From the generating function of F_{n+2} , we obtain

$$\sum_{n \geq 2} n F_{n+2} x^n = x \left(\frac{x^2(3+2x)}{1-x-x^2} \right)' = \frac{x^2(6+3x-4x^2-2x^3)}{(1-x-x^2)^2},$$

which yields that

$$(1-x)u(x) = x^2 \left[\frac{x^2(3+2x)}{1-x-x^2} + \frac{x^2(6+3x-4x^2-2x^3)}{(1-x-x^2)^2} + u(x) \right] + x^2 + 4x^3.$$

Therefore,

$$u(x) = x^2(1+2x)/(1-x-x^2)^3,$$

which gives the generating function for $f_{312}(n)$ as shown in formula (3.3).

For formula (3.4), we first have

$$\sum_{n \geq 3} f_{321}(n) x^n = \sum_{n \geq 3} \binom{n}{3} F_{n+1} x^n - 2 \sum_{n \geq 3} f_{312}(n) x^n \quad (3.7)$$

from $2f_{312}(n) + f_{321}(n) = \binom{n}{3} F_{n+1}$. Using the fact that $\sum_{n \geq 0} F_n x^n = \frac{x}{1-x-x^2}$, we get

$$\begin{aligned} \sum_{n \geq 3} F_{n+1} x^n &= \frac{1}{x} \left(\frac{x}{1-x-x^2} - x - x^2 - 2x^3 \right) = \frac{x^3(3+2x)}{1-x-x^2}, \\ \sum_{n \geq 3} \binom{n}{3} F_{n+1} x^n &= \frac{x^3}{6} \left(\sum_{n \geq 3} F_{n+1} x^n \right)''' = \frac{x^3(3+8x+6x^2+4x^3)}{(1-x-x^2)^4}. \end{aligned}$$

Formula (3.4) follows by substituting the generating functions of $\binom{n}{3} F_{n+1}$ and $f_{312}(n)$ into (3.7), and we complete the proof. $\blacksquare$

The first few values of $f_q(S_n(123, 132, 213))$ for q of length 3 are shown below.

n	f_{123}	f_{132}	f_{213}	f_{231}	f_{312}	f_{321}	n	f_{123}	f_{132}	f_{213}	f_{231}	f_{312}	f_{321}
3	0	0	0	1	1	1	6	0	0	0	40	40	180
4	0	0	0	5	5	10	7	0	0	0	95	95	545
5	0	0	0	15	15	50	8	0	0	0	213	213	1478

3.2 Pattern Count on Other Triple Avoiding Permutations

We begin with the pattern count on $(123, 132, 231)$ -avoiding permutations.

Theorem 3.5. *For $n \geq 3$, in the set $S_n(123, 132, 231)$, we have*

$$f_{213}(n) = f_{312}(n) = \binom{n}{3}, \quad (3.8)$$

$$f_{321}(n) = (n-2) \binom{n}{3}. \quad (3.9)$$

Proof. We first give the following structure from Simion and Schmidt [8]

$$\sigma \in S_n(123, 132, 231) \Leftrightarrow \sigma = n, n-1, \dots, k+1, k-1, k-2, \dots, 1, k \text{ for some } k. \quad (3.10)$$

Based on such structure, we can show $f_{213}(n) = f_{312}(n)$ by a direct bijection. Let $q = abc$ be a 213-pattern of a permutation $\sigma \in S_n(123, 132, 231)$. From $b < c$ and the fact that $\sigma \in S_n(123, 132, 231)$ has only one ascent at position $n-1$, we have $\sigma(n) = c$, and thus $\sigma = n, n-1, \dots, c+1, c-1, c-2, \dots, 2, 1, c$ from the structure (3.10). Let $q' = cba$ (a 312-pattern) and we set $\sigma' = n, n-1, \dots, a+1, a-1, a-2, \dots, 2, 1, a$ as the desired permutation. For example, if $n = 7$ and $q = 326$, then $\sigma = 7543216$. Moreover, $q' = 623$ and $\sigma' = 7654213$.

To calculate $f_{312}(n)$, we suppose $\sigma = n, n-1, \dots, k+1, k-1, k-2, \dots, 2, 1, k$ for some k from structure (3.10). We can construct a 312-pattern as follows: Choose one element from the first $n-k$ elements to play the role of 3, then choose one element from the next $k-1$ elements to play the role of 1, and the last element plays the role of 2. Thus, summing up all possible k , we have

$$f_{312}(n) = \sum_{k=1}^n (n-k)(k-1) = -n^2 + (n+1) \sum_{k=1}^n k - \sum_{k=1}^n k^2 = \frac{n(n-1)(n-2)}{6} = \binom{n}{3}.$$

The formula for $f_{321}(n)$ follows by using $f_{213}(n) + f_{312}(n) + f_{321}(n) = n \binom{n}{3}$. $\blacksquare$

The first few values of $f_q(S_n(123, 132, 231))$ for q of length 3 are shown below.

n	f_{123}	f_{132}	f_{213}	f_{231}	f_{312}	f_{321}	n	f_{123}	f_{132}	f_{213}	f_{231}	f_{312}	f_{321}
3	0	0	1	0	1	1	6	0	0	20	0	20	80
4	0	0	4	0	4	8	7	0	0	35	0	35	175
5	0	0	10	0	10	30	8	0	0	56	0	56	336

For $(132, 213, 231)$ -avoiding permutations, we have

Theorem 3.6. *For $n \geq 3$, in the set $S_n(132, 213, 231)$,*

$$f_{123}(n) = f_{312}(n) = \binom{n+1}{4}, \quad (3.11)$$

$$f_{321}(n) = \frac{n(n-2)(n-1)^2}{12}. \quad (3.12)$$

Proof. We begin with the following structure by Simion and Schmidt [8]

$$\sigma \in S_n(132, 213, 231) \Leftrightarrow \sigma = n, n-1, \dots, k+1, 1, 2, \dots, k-1, k \text{ for some } k. \quad (3.13)$$

Based on such structure, we first prove $f_{123}(n) = f_{312}(n)$. For each

$$\sigma = n, n-1, \dots, k+1, 1, \dots, \underline{a}, a+1, \dots, \underline{b}, b+1, \dots, c-1, \underline{c}, c+1, \dots, k-1, k$$

with abc as a 123-pattern, we set

$$\sigma' = n, n-1, \dots, \underline{n-k+c}, \dots, c, 1, 2, \dots, \underline{a}, a+1, \dots, \underline{b}, b+1, \dots, c-1,$$

and it is easy to check that $n-k+c, a, b$ is a 312-pattern of σ' . For example, if $\sigma = 98712\underline{3}4\underline{5}6$ then $\sigma' = 9\underline{8}7651\underline{2}34$.

To calculate $f_{123}(n)$, we suppose $\sigma = n, n-1, \dots, k+1, 1, 2, \dots, k-1, k$ for some k from structure (3.13). Then, a 123-pattern can be obtained by choosing three elements from the last k elements to play the role of 123. Thus, summing up all possible k gives

$$f_{123}(n) = \sum_{k=1}^n \binom{k}{3} = \binom{n+1}{4}.$$

We complete the proof by using $f_{123}(n) + f_{312}(n) + f_{321}(n) = n \binom{n}{3}$. ■

The first few values of $f_q(S_n(132, 213, 231))$ for q of length 3 are shown below.

n	f_{123}	f_{132}	f_{213}	f_{231}	f_{312}	f_{321}	n	f_{123}	f_{132}	f_{213}	f_{231}	f_{312}	f_{321}
3	1	0	0	0	1	1	6	35	0	0	0	35	50
4	5	0	0	0	5	6	7	70	0	0	0	70	105
5	15	0	0	0	15	20	8	126	0	0	0	126	196

For (123, 132, 312)-avoiding permutations, we have

Theorem 3.7. For $n \geq 3$, in the set $S_n(123, 132, 312)$,

$$f_{213}(n) = f_{231}(n) = \binom{n}{3}, \quad (3.14)$$

$$f_{321}(n) = (n-2) \binom{n}{3}. \quad (3.15)$$

Proof. We begin with the following structure from Simion and Schmidt [8]

$$\sigma \in S_n(132, 213, 231) \Leftrightarrow \sigma = n-1, n-2, \dots, k+1, n, k, k-1, \dots, 1 \text{ for some } k. \quad (3.16)$$

Based on such structure, we prove $f_{213}(n) = f_{231}(n)$ by a direct correspondence. Let $\sigma = n-1, \dots, \underline{a}, a+1, \dots, \underline{b}, b+1, \dots, k+1, \underline{n}, k, k-1, \dots, 2, 1$ with abn as a 213-pattern. Then, we set

$$\sigma' = n-1, \dots, \underline{n-a+b}, \dots, n-a+k+1, \underline{n}, n-a+k, n-a+k-1, \dots, \underline{n-a}, \dots, 2, 1,$$

where $n-a+b, n, n-a$ is a 231-pattern of σ' . For example, if $\sigma = 8 \underline{7} 6 \underline{5} 4 \underline{9} 3 2 1$, then $\sigma' = \sigma = 8 \underline{7} 6 \underline{9} 5 4 \underline{3} 2 1$.

To calculate $f_{213}(n)$, we suppose that $\sigma = n-1, n-2, \dots, k+1, n, k, k-1, \dots, 2, 1$ for some k . Then, a 213-pattern can be obtained by choosing two elements from the first $n-k-1$ elements to play the role of 21, and let n play the role of 3. Thus, summing up all possible k , we have

$$f_{213}(n) = \sum_{k=0}^{n-1} \binom{n-k-1}{2} = \binom{n}{3}.$$

We complete the proof by using $f_{213}(n) + f_{231}(n) + f_{321}(n) = n \binom{n}{3}$. ■

The first few values of $f_q(S_n(123, 132, 312))$ for q of length 3 are shown below.

n	f_{123}	f_{132}	f_{213}	f_{231}	f_{312}	f_{321}	n	f_{123}	f_{132}	f_{213}	f_{231}	f_{312}	f_{321}
3	0	0	1	1	0	1	6	0	0	20	20	0	80
4	0	0	4	4	0	8	7	0	0	35	35	0	175
5	0	0	10	10	0	30	8	0	0	56	56	0	336

Finally, we study the pattern count on $(123, 231, 312)$ -avoiding permutations.

Theorem 3.8. *For $n \geq 3$, in the set $S_n(123, 231, 312)$, we have*

$$f_{132}(n) = f_{213}(n) = \binom{n+1}{4}, \quad (3.17)$$

$$f_{321}(n) = \frac{n(n-2)(n-1)^2}{12}. \quad (3.18)$$

Proof. From Lemma 2.1, we see that

$$\sigma \in S_n(123, 231, 312) \Leftrightarrow \sigma^r \in S_n(321, 132, 213) \Leftrightarrow (\sigma^r)^c \in S_n(123, 231, 312).$$

We have $f_{213}(n) = f_{132}(n)$ from $(213^r)^c = 312^c = 132$. The following structure of the set $S_n(123, 231, 312)$ is given by Simion and Schmidt [8]

$$\sigma \in S_n(132, 213, 231) \Leftrightarrow \sigma = k-1, k-2, \dots, 3, 2, 1, n, n-1, \dots, k \text{ for some } k. \quad (3.19)$$

Suppose that $\sigma = k-1, k-2, \dots, 3, 2, 1, n, n-1, \dots, k$ for some k , then a 213-pattern can be obtained as follows: Choose two elements from the first $k-1$ elements to play

the role of 21, and choose one element from the last $n - k + 1$ elements to play the role of 3. Thus, summing up all possible k , we have

$$f_{213}(n) = \sum_{k=1}^n \binom{k-1}{2} (n - k + 1) = \sum_{k=0}^{n-1} \binom{k}{2} (n - k) = \binom{n+1}{4}.$$

The formula for $f_{321}(n)$ is obtained by the relation $2f_{213}(n) + f_{321}(n) = n \binom{n}{3}$. ■

The first few values of $f_q(S_n(123, 231, 312))$ for q of length 3 are shown below.

n	f_{123}	f_{132}	f_{213}	f_{231}	f_{312}	f_{321}	n	f_{123}	f_{132}	f_{213}	f_{231}	f_{312}	f_{321}
3	0	1	1	0	0	1	6	0	35	35	0	0	50
4	0	5	5	0	0	6	7	0	70	70	0	0	105
5	0	15	15	0	0	20	8	0	126	126	0	0	196

References

- [1] M. Bóna, A Walk Through Combinatorics, 3rd edition, World Scientific, 2011.
- [2] M. Bóna, Surprising symmetries in objects counted by Catalan numbers, Electron. J. Combin. 19 (2012), P62.
- [3] M. Bóna, The absence of a pattern and the occurrences of another, Discrete Math. Theor. Comput. Sci. 12 (2010), no. 2, 89–102.
- [4] L. Comtet, Advanced Combinatorics, Reidel, Dordrecht, 1974.
- [5] J. Cooper, Combinatorial Problems I like, internet resource, available at <http://www.math.sc.edu/~cooper/combprob.html>.
- [6] C. Homberger, Expected Patterns in Permutation Classes, Electron. J. Combin. 19(3) (2012), P43.
- [7] K. Rudolph, Pattern Popularity in 132-avoiding Permutations, Electron. J. Combin. 20(1) (2013), P8.
- [8] R. Simion and F.W. Schmidt, Restricted permutations, European J. Combin. 6 (1985), 383–406.
- [9] R.P. Stanley, Enumerative Combinatorics, vol. 1, Cambridge University Press, 1997.